

The Relationship between Eating Disorders and ICAM-1, E-selection and Ghrelin Resting Level in Overweight People

*Gholamreza Sharifi¹

¹Assistant Professor of Sports Science, Islamic Azad University, Khorasgan Branch, Isfahan, Iran.

Abstract

Introduction

There is an agreement that eating disorder is related to psychological characteristics and on the other hand, level of ghrelin hormone, Intercellular adhesion molecule-1 (ICAM-1) and E-selection also change during eating disorders. We aimed to study the relationship between eating disorders and rest levels, ICAM-1, E-selection, and ghrelin hormone in obese people.

Materials and Methods

In this quasi-experimental study, 120 obese men (25-30 years old) were purposefully selected. Then the data about their eating disorders gathered with eating attitudes test (EAT-26) questionnaire. In the next phase in the rest condition and after overnight fasting, blood samples are collected for measurement of rest levels, ICAM-1, E-selection, and ghrelin hormone. Finally the data were analyzed with appropriate statistical tests in SPSS version 18.

Results

Mean and deviation of rest levels, ICAM-1, E-selection, and ghrelin hormone were respectively 3064.19, 61.5±19.7, and 2.5±1.5 and there was not any statistical significance relationship between eating disorders ICAM-1, E-selection, and ghrelin hormone in obese men ($P<0.05$).

Conclusion

According to our results, the prevalence of eating disorders can be a reason other than rest Levels, ICAM-1, E-selection, and ghrelin hormone.

Keywords: Eating disorders, E-selection, Ghrelin, ICAM-1.

***Corresponding Author:**

Dr. Gholamreza Sharifi, Islamic Azad University, Khorasgan Branch, Isfahan, Iran.

Email: gre_sharifi@yahoo.com

Received date: Jun 24, 2014 ; Accepted date: Jul 22, 2014

Introduction

After obesity and asthma, eating disorders are the third most common cause of disease in the young people of societies (1-3). There is a significant relation between ghrelin levels and nutritional conditions and energy balance, so that some researchers believe that ghrelin effects on increasing of fat storage should be taken independently from its effort on growth hormone (4).

Weight control, permanent mental concern of food and dissatisfaction from body image understanding, lead to some sort of nutritional - psychological disorders that is known as eating disorders. The main and obvious features of these disorders are use of different kind of diets and weight control methods, disorganization of balanced nutritional behaviors and causing evident thought and individual imaginations disorders concerning the food and its own (specially about weight). These disorders have numerous nutritional and psychological effects and may cause to some problems, such as stress and depression and if they would not diagnose it may lead to intense and clinical cases as bulimia nervosa and anorexia nervosa (5).

The prevalence of obesity and overweight is dangerously increasing through the world. There is also a high present of obesity and overweight probably in different sex and age groups in our country (6). Obesity and overweighting are of the most important unavoidable factors that may lead to illness and death.

There are some sticky molecules which have important effect on cardiovascular diseases and specially atherosclerosis and mostly in overweight people, including ICAM-1 and E- selection and ghrelin. These molecules are glycoprotein receptors that lie on the outer surface of the cell membrane and reach to cytoplasm after crossing cell membrane. They motive

white blood cells to movement to lymphoid tissues and especially to infectious and inflamed areas and take part in vascularization, wound healing, inflammation, and development and coordination of vessels processes(6). Previous researches have shown that ghrelin hormone has strong effects in commencement of getting the food so that by increasing of this hormone man feels hungry. Establishment secretion of ghrelin in stomach and gastric – intestinal rout, ghrelin receptor synthesis in Valgus nerve and numerous gastric – intestinal effects of this hormone, indicate the significant role of ghrelin in brain stem gastric (7,8).The advantage of ghrelin as hunger hormone with its effects on appetite that is created by hypothalamic peptides, Neuropeptide Y (NPY), and Agouti-related protein (AGRP).

According to continual concerns of overweight people, in this type of diseased, the probability of eating disorders is higher than others and the prevalence of these disorders can be an important step in nutrition planning in order to prevention and treatment and may have effective role in their health promotion. There are also some investigations subject to ICAM-1 and E-selection resting levels and ghrelin in obese people, such as Boos (2008), Mensud (2007), Josef (2005), Dace (2012)(9-12). Therefore this study intended to examined the relation between eating disorders and ICAM-1 and E-selection resting levels and ghrelin in overweight men.

Materials and Methods

In this study we invite 120 overweight 30-35 years old men who had come into Alzahra Hospital in Isfahan, the capital of Isfahan Province. Then they showed their consent to participated in the study. They were non smokers and had no diabetes or endocrine and disease. They were

explained about conditions then completed the testimonial forms and participate in this research. In this study the result of hormonal evaluation kept confidential and after examining the result, their hormonal evaluations were given to them. This study has accomplished in multistage thus that after filling the questionnaire, in sitting mode and fasting condition, 5 cc venous blood, was taken from left hand's antecubital vein. After 5 minutes for coagulation, the mentioned samples were centrifuged with 3000 Revolutions per minute (RPM) for 10 minutes. Blood tests have carried out to determining the serum ghrelin levels, ICAM and E-selection. The tribals were performed by quantitative detection kit, under the care of the researcher in the specialized laboratory. Then each group eating disorders data collected through 26 question questionnaire.

Afterward, in resting condition and after overnight fasting, a blood sample was taken and after separating its plasma, it was kept in -20 degree in laboratory until the ghrelin resting levels, ICAM and E-selection measuring time.

Table1. The relation between eating disorders and ghrelin, ICAM and E-Selection resting levels in overweight people

Variables	Numbers	Mean \pm SD	Eating disorders	Pearson correlation coefficient	P value
Ghrelin	120	3064.19 \pm 371.61	0.8	0.044	0.816
ICAM	120	61.5 \pm 19.7	0.8	0.06	0.489
E- Selection	120	2.5 \pm 1.5	0.8	-0.094	0.278

Discussion

In the present study the prevalence of eating disorders relative to ghrelin, ICAM and E-selection resting levels in overweight men was determined, but no significant relation was founded. Eating disorders is under influence of several factors such as social – psychological

The data analysis is performed by spss18 software and according to the data normality, the analyzing of the data is done by descriptive statistics and inferential statistics. For examining the normality of the data Kolmogorov-Smirnov test was applied and to assess the relation between variables, t-test and Pearson correlation coefficient and correlation test was used. Additionally the level of confidence for all the tests was considered to 95%.

Results

Based on the results of investigations Body mass index (BMI) of the participants was: 25 – 30 Kg/m², weight range was: 87.4 \pm 9.3 and age group was: 25 – 30 years old. As it shown the average and standard deviation of the ghrelin resting levels, ICAM and E-selection were 3064.19 \pm 371.61, 61.5 \pm 19.7, 2.5 \pm 1.5, respectively.

Therefore there was no significant relation between eating disorders Pearson correlation coefficient and ghrelin, ICAM and E-selection hormones (P> 0.05).

mechanisms, genetics and physiology. Eating problems and weight control behaviors, cover apply to wide range of people. In the last decades a considerable growth in prevalence of weight disorders and eating problems in obese people is observed. In the most of studies of ghrelin basic levels in various types of clinical eating disorders, such as anorexia nervosa

and bulimia nervosa is reported more than basic level of them in healthy people (10-12). Some investigations result indicated that in medical conditions and reaching to the normal weight ghrelin basic levels will decrease and in some cases they will decrease more than the level of this hormone in healthy people (13). The results of this research subject to ICAM-1 and E-selection serum level in comply with Anki Tonjes (2007), Saetre (2011), Hejazi (2013), Robert (2006), Wang (2001), Sampson (2006), Adamopoulos (2001), Rothenbacher (2003), Zoppini (2006) (17-26).

It is reported that inflammation and notice to Cardiovascular and Medical Sciences (CAMs) as cardiovascular risk factors is very important in people whit coronary heart disease. These factors have strong relations with the other risk factors such as CRP, cholesterol and lipoprotein A. Endomyocardial represent the considerable relation between increasing of CRP and ICAM-1 in vascular endothelial. And also the interaction between lipoprotein A and ICAM-1 is indicated (27). In inflammation condition by increasing the circulatory system activity that is one of the important parts of inflammation responses, vessels endothelial damages will be extended and the molecules which are commencer in topical delivery of leukocytes will be active. Sticky molecules, such as ICAM-1, can be released (specific or nonspecific) by inflamed tissue damage, for example nonspecific proteolysis (28). Regular exercise by decreasing the sympathetic stimulation and increasing the anti-inflammatory cytokine can restrain the releasing of Interleukin 1 beta (IL-1 β) and Tumor necrosis factor alpha (TNF- α) pro-inflammatory cytokines from adipose tissue and consequently the serum concentration of ICAM-1 and E-selection will decrease (29). Doing exercise by decreasing harmful lipolysis and

increasing blood helpful lipolysis, decrease the risk of cardiovascular diseases.

The endocrine system, can increase lipid oxidation (lipolysis) by accretion epinephrine, norepinephrine and growth hormones and cortisol and therefore provide the required energy for the muscles. During these actions, after 30 to 45 minutes of activity commencement, plasma cortisol value reach to maximum and by increasing the gluconeogenesis process, accelerate calling and using of free fatty acids for producing energy during the exercise (30). Since adipose tissue is one of the important areas for secretion of inflammatory markers and cytokines, endurance exercises by increasing the lipolysis and decreasing the body fat can be a mechanism for reducing the inflammatory mediators and intercellular sticky molecules.

Conclusion

The results of this study indicated that the eating disorders prevalence can happen in consequence of reasons other than ghrelin, ICAM-1 and E-selection resting levels, and according to the determined percent of eating disorders prevalence in overweight people, there would be a need to inform people about the healthy and unhealthy methods of weight control and proper nutrition.

Conflict of interests: None

Acknowledgment

Special thanks to this research on tribals for their sincere cooperation. This study complied with the current laws of Iran. For preparing blood samples from subjects, consent form completed by them.

References

1. Ljubuncic P, Portnaya I, Cogan U, Azaizeh H, BomzonA. Antioxidant activity of *Crataegus aronia* aqueousextract used in traditional Arab

- medicine in Israel. *J Ethnopharmacol* 2005; 101: 153-61.
2. Khodabakhsh MR, Kiani F. Effects of Emotional Eating on Eating Behaviors Disorder in Students: The Effects of Anxious Mood and Emotion Expression. *International J of Pediatrics* 2014; 2(4.1): 295-303.
 3. Kiani F, Khodabakhsh MR. Perfectionism and Stressful Life Events as Vulnerabilities to Depression Symptoms in Students. *International J of Pediatrics* 2014; 2(4.1): 277-285.
 4. Hansen TK, Dall R, Hosoda H, Kojima M, Kangawa K, Vhristiansen JS, et al. Weight loss increases circulating levels of ghrelin in human obesity. *Journal of clinical endocrinology* 2002;56:203-6.
 5. O'Brien KA, Ling S, Abbas E, Dai A, Zhang J, Wang WC, et al. A Chinese Herbal Preparation Containing Radix Salviae Miltiorrhizae, Radix Notoginseng and Borneolum Reduces Circulating Adhesion Molecules. *Evid Based Complement Alternat Med* 2011;2011:790784.
 6. Bayati M. Physical inactivity and sedentary lifestyle. *Iran J Endocrinol Metabol* 2012; 13(5): 537-9.
 7. Mogharnasi M, Gaeini AA, Sheikholeslami Vatani D. Effect of sprint training and detraining period on cellular adhesion molecule (sICAM-1) in wistar rats. *Olympic* 2008; 16:19-30.
 8. Wu JT, Kral JG. Ghrelin integrative neuro endocrine peptide in health and disease. *Annals of surgery* 2004; 239: 464-74.
 9. Boos CJ, Balakrishnan B, Lip GY. The effects of exercise stress testing on soluble E-selectin, von Willebrand factor, and circulating endothelial cells as indices of endothelial damage/dysfunction. *Ann med* 2008;40(1):66-73.
 10. Mensud H. Vascular inflammatory markers in early-onset obese and type 2 diabetes subjects before and after three months' aerobic exercise training. *Diabetes and Vascular Disease Research* 2007;4: 231.
 11. Josef N. Exercise training in chronic heart failure: effects on pro-inflammatory markers. *The European Journal of Heart Failure* 2005; 7 : 189-93.
 12. Dace R. Systemic pro-inflammatory molecule response to acute submaximal exercise in moderately and highly trained athletes. *Environmental and Experimental Biology* 2012; 10: 107-112.
 13. Prine A C, Brooks S J, Stahl D, Treasure J. Systematic review and meta analysis of the baseline concentration and physiologic responses of gut hormones to food in eating disorders. *American society for nutrition* 2009; 89(3): 755-65.
 14. Troisi A, Lorenzo G, Lega I, Tesaro M, Bertoli A, Leo R, Iantorno M, et al. Plasma ghrelin in anorexia, bulimia, and binge-eating disorder: relations with eating patterns and circulating concentrations of cortisol and thyroid hormones. *Neuroendocrinology* 2005;81:259-66.
 15. Geleibter A, Gluck ME, Hashim SA. Plasma ghrelin concentrations are lower in binge-eating disorder. *Journal nutrition* 2005;135: 1326-30.
 16. Janas -kozik M, Krupka-matuszczyk I, Malinowska -kolodziej I, Lewin-kowalik J. Total ghrelin plasma level in patients with the restrictive type of anorexia nervosa. *Journal of regulatory peptides* 2007;140: 4346.
 17. Tönjes A, Scholz M, Fasshauer M, Kratzsch J, Rassoul F, Stumvoll M, et al. Beneficial effects of a 4-week exercise program on plasma concentrations of adhesion molecules. *Diabetes Care* 2007 Mar; 30(3):e1.
 18. Saetre T, Enoksen E, Lyberg T, Strandén E, Jørgensen JJ, Sundhagen JO, et al. Supervised exercise training reduces plasma levels of the endothelial inflammatory markers E-selectin and ICAM-I in patients with peripheral arterial disease. *Angiology* 2011;62(4):301-5.
 19. Hejazi M, Hosseini Abrishami L, Mohammad Khani J, Boghrabadi V. The Effects of 8-Week Aerobic Exercises on Serum Levels of Cell Adhesion Molecules among Middle-Aged Women. *Advanced Studies in Biology* 2013;. 5(6): 279 -89.
 20. Roberts CK, Won D, Pruthi S, Lin SS, Barnard RJ. "Effect of a diet and exercise intervention on oxidative

stress, inflammation and monocyte adhesion in diabetic men". *Diabetes Res Clin Pract* 2006; 73(3):249-59.

21. Wang RY, Yang YR, Yu SM. "Protective effects of treadmill training on infarction in rats". *Brain Res* 2001; 922(1):140-3.
22. Ding YH, Young CN, Luan X, Li J, Rafols JA, Clark JC, et al. "Exercise preconditioning ameliorates inflammatory injury in ischemic rats during reperfusion". *Acta Neuropathol* 2005; 109(3):237-46.
23. Richard J, Simpson, Geraint D, Florida-James, Greg P, Whyte, Keith Guy. "The effects of intensive, moderate and downhill treadmill running of human blood lymphocytes expression the adhesion activation molecules CD54 (ICAM-1), CD18 (B2 integrin) and CD53". *European Journal of Applied Physiology* 2006; 97(1): 109-121.
24. Adamopoulos S, Parissis J, Kroupis C, Georgiadis M, Karatzas D, Karavolias G, et al. "Physical training reduces peripheral marker of inflammation in patients with chronic heart failure". *European Heart Journal* 2001; 22, 791-97.
25. Rothenbacher D, Hoffmeister A, Brenner H, Koenig W. Physical activity, coronary heart disease, and Inflammatory response. *Archives of Internal Medicine* 2003; 163: 1200-1205.
26. Zoppini G, Targher G, Zamboni C, Venturi C, Cacciatori V, Moghetti P, et al. Effects of moderate-intensity exercise training on plasma biomarkers of inflammation and endothelial dysfunction in order patients with type 2 diabetes. *Nutrition, Metabolism and Cardiovascular Diseases* 2006; 16: 543-9.
27. Davies MJ, Gordon JL, Gearing AJ, Pigott R, Woolf N, Katz D, et al. Circulating Adhesion Molecules in a Ten-year Follow-up Study After Premature Myocardial Infarction. *The Open Inflammation Journal* 2008; 171(3):223-9.
28. Macías C, Villaescusa R, del Valle L, Boffil V, Cordero G, Hernández A, et al. Endothelial Adhesion Molecules ICAM-1, VCAM-1 and E-selectin in Patients With Acute Coronary Syndrome. *Rev Esp Cardiol* 2003 Feb; 56(2):137-44.
29. Mohammad-Ali V, Bulmer K, Clarke D, Goodrick S, Coppack SW, Pinkney JH. Adrenergic regulation of proinflammatory cytokines in humans. *International Journal of obesity and Related Metabolic Disorders* 2000; 24: 154-5.
30. Wegge JK, Roberts CK, Ngo T, Bamard RJ. Effect of diet and exercise intervention on inflammatory and adhesion molecules in postmenopausal women on hormone. *Metabolism* 2004; 53(3):377-81.