

Infant and Young Child Feeding Status in Iran Compared the Different United Nation Regions

Salman Khazaei¹, Kamyar Mansori², Zaher Khazaei³, Mohadeseh Sani⁴, *Erfan Ayubi⁵

¹Department of Epidemiology, School of Public Health, Hamadan University of Medical Sciences, Hamadan, Iran. ²Social Development & Health Promotion Research Center, Gonabad University of Medical Sciences, Gonabad, Iran. ³Social Determinants of Health Research Center, Kurdistan University of Medical Sciences, Sanandaj, Iran. ⁴School of Medicine, Zabol University of Medical Sciences, Zabol, Iran. ⁵Department of Epidemiology, School of Public Health, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

Dear Editor-in-Chief,

Promote and support breastfeeding is an essential element for the health and development of infants so that, it introduced as most cost-effective intervention to reduce infant mortality in developing countries (1). Breastfeeding has an important role to ensure food security for maintain a healthy and productive life for a large proportion of babies in the world, in addition, it is associated with improves intelligence quotient (IQ), school attendance and higher income in adult life (2).

World Health Organization (WHO) indicated exclusive breastfeeding for the first six months of life with starting in the first half hour after delivery for optimal growth and development (3). Early initiation of breastfeeding (first hour after birth) has positive effects on biological and emotional health for mother and child and reduces child mortality (4). It is estimated that if breastfeeding in children aged at 0-23 months have been considered optimally, among children under 5 years old globally over 800,000 life could be saved every year (2). In other hand, protect effects of breastfeeding for mothers against breast and ovarian cancer, osteoporosis have been identified (5).

Low rates and early cessation of breastfeeding can be along with important adverse effects on health. In this letter, we aimed to describe the profile of breastfeeding in Iran compared with other regions of the world, and to express some solutions for promotion of breastfeeding in regard to published studies about this matter.

Key Words: Breastfeeding, Exclusive breast feeding, Children, Iran, United Nation Regions.

Please cite this article as: Khazaei S, Mansori K, Khazaei Z, Sani M, Ayubi E. Infant and Young Child Feeding Status in Iran Compared Different United Nation Regions. *Int J Pediatr* 2016;4(10):3639-41. DOI: **10.22038/ijp.2016.7396**

*Corresponding Author:

Erfan Ayubi (MSc, PhDc), Department of Epidemiology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran.

Email: aubi65@gmail.com

Received date Jul 20, 2016; Accepted date: Aug 22, 2016




Fig.1: Globally percent of infants < 6 months of age exclusively breastfed in 2015

Table-1: Infant and young child feeding status in Iran compared different UN regions

Regions	Early initiation of breastfeeding (<1 hour, %)	Exclusive breastfeeding (<6 months, %)	Breastfeeding at 1 year (12–15 months, %)
Africa	46	40	86
Asia	41	40	71
Latin America and the Caribbean	49	32	51
Central and Eastern Europe	54	27	56
Iran	69	53	84
World	44	39	74

As illustrated in **Figure.1** and **Table.1** (4), Africa and Southern East Asia have lower rate of exclusive breastfeeding in children under 6 month and early initiation of breastfeeding under one hour. In Iran, rate of two former indicators has higher than other part of world and the estimated global averages. Higher percentage of breastfeeding at one year was observed in Africa and Asia. In Iran percentage of breastfeeding at one year was 84% (4). In Iran compared other regions, early initiation of breastfeeding, exclusive breastfeeding and breastfeeding at one year have suitable rates; but it should be noted that about 50% of children have

early cessation of breastfeeding and the results were far from meeting the WHO recommendation (6). Some national studies indicated that considering breastfeeding in the first hour after birth can be associated with mother's educational level, gestational age, previous history of breastfeeding, vaginal delivery, infant weight, infant disease, hospitalization in Neonatal intensive care unit (NICU), rooming in and delivery in public hospitals (7). Enhancing the knowledge of mothers about importance of breastfeeding in the first 6 months of baby's life is undeniable (8). Physician's recommendation, insufficient breast milk,

maternal or infant illness and working mother are some of other causes (9). With considering and embedding some strategies into health care programs, exclusive breastfeeding can be achievable. These strategies are as follow;

1) Breastfeeding policy and/or action plans, 2) Leadership for breastfeeding promotion, 3) Training of health care staff, 4) Baby Friendly Hospital Initiative, 5) Endorsement of the WHO International Code of Marketing for breast milk substitutes, 6) Legislation for working mothers, 7) Community outreach, including mother support, 8) Information, education, communication, 9) Monitoring of breastfeeding duration and exclusiveness and 10) Activities directed towards disadvantaged groups (10).

Conclusion

Estimation of the mentioned indicators is essential to guide investment to improve nutrition and health during the first two years of life. Although, compared other UN regions, Iran has a good situation in this regard, but it is necessary the feeding status rates to be enhanced. The factors which affect these indices are various and complex, thus to achieve adequate and sustained increases in breastfeeding, many other programs including: employment legislation, widespread public education, community support and improved health care practices are needed.

REFERENCES

1. Fjeld E, Siziya S, Katepa-Bwalya M, Kankasa C, Moland KM, Tylleskär T. 'No sister, the breast alone is not enough for my baby'a qualitative assessment of potentials and barriers in the promotion of exclusive breastfeeding in southern Zambia. *International breastfeeding journal*. 2008;3(26):DOI: 10.1186/746-4358-3-26.
2. Victora CG, Bahl R, Barros AJ, França GV, Horton S, Krasevec J, et al. Breastfeeding in the 21st century: epidemiology, mechanisms, and lifelong effect. *The Lancet*. 2016;387(10017):475-90.
3. Kramer MS, Kakuma R. The optimal duration of exclusive breastfeeding. *Protecting Infants through Human Milk*: Springer; 2004. p. 63-77.
4. UNICEF. Infant and Young Child Feeding database. 2016 [updated Aug 2016; cited Jul 2016]; Available from: <http://data.unicef.org/nutrition/iycf.html>.
5. Ball O. Breastmilk is a human right. *Breastfeeding Review*. 2010;18(3):9-19.
6. World Health Organization. The World Health Organization's infant feeding recommendation. 2016 [cited 21 Sep 2016]; Available at: http://www.who.int/nutrition/topics/infantfeeding_recommendation/en/.
7. Haghghi M, Taheri E. Factors Associated with Breastfeeding in the First Hour after Birth, in Baby Friendly Hospitals, Shiraz-Iran. *International Journal of Pediatrics*. 2015;3(5.1):889-96.
8. Esfandtari R, Baghiani Moghadam MH, Khakshour A, Faroughi F, Zarif B, Saeidi M. Study of Maternal Knowledge and Attitude toward Exclusive Breast Milk Feeding (BMF) in the First 6 Months of Infant in Yazd-Iran. *International Journal of Pediatrics*. 2014;2(3.1):175-81.
9. Olang B, Heidarzadeh A, Strandvik B, Yngve A. Reasons given by mothers for discontinuing breastfeeding in Iran. *International breastfeeding journal*. 2012;7(1):DOI: 10.1186/746-4358-7-7.
10. Olang B, Farivar K, Heidarzadeh A, Strandvik B, Yngve A. Breastfeeding in Iran: prevalence, duration and current recommendations. *International breastfeeding journal*. 2009;4(1):DOI: 10.1186/746-4358-4-8.