

The Correlation of Leukocyte Count and Percentage of Segmented Neutrophils with Pathohistological Findings of Appendix in Children

*Marko Baskovic¹, Bozidar Zupancic², Ana Dominkovic³, Ante Cizmic⁴, Davor Jezek⁵

¹Department of Pediatric Surgery, Children's Hospital Zagreb, Klaićeva 16, Zagreb 10000, Croatia.

²Professor, Department of Pediatric Surgery, Children's Hospital Zagreb, Klaićeva 16, Zagreb 10000, Croatia.

³University of Zagreb, School of Medicine, Šalata 3, Zagreb 10000, Croatia.

⁴Assistant professor, Department of Pediatric Surgery, Children's Hospital Zagreb, Klaićeva 16, Zagreb 10000, Croatia.

⁵Professor, Department of Histology and Embriology, University of Zagreb, School of Medicine, Šalata 3, Zagreb 10000, Croatia.

Abstract

Background: Appendicitis is the most common indication for an emergency operation in children's age. Although none of the laboratory values has not high sensitivity and specificity for the diagnosis of appendicitis, leukocyte count and the percentage of segmented neutrophils are most commonly used. The aim of this study was to determine whether there is a statistically significant correlation between leukocyte count and the percentage of segmented neutrophils compared to the pathohistological finding of appendix in children.

Materials and Methods: We retrospectively analyzed the data in the period from 1 January 2016 to 31 December 2016. The analysis was made on 211 patients (from the Children's Hospital Zagreb, Croatia). Spearman's correlation coefficient (r_s) was calculated. We determined the specificity and sensitivity of leukocyte count and the percentage of segmented neutrophils used in the calculation of Alvorado and Pediatric Appendicitis score.

Results: The results of the research have shown that the correlation between leukocyte count and the pathohistological findings is weak ($r_s = 0.29$, $p = 3.61 \cdot 10^{-8}$); while there is no correlation between the percentage of segmented neutrophils and pathohistological findings ($r_s = 0.18$, $p = 7.08 \cdot 10^{-5}$). The sensitivity of leukocyte count is 93% and the specificity is 30%, while the sensitivity to the percentage of segmented neutrophils is 71% and the specificity is 50%. The Receiver Operating Characteristic (ROC) curve analysis for leukocytes shows area under the curve of 0.648, while for segmented neutrophils of 0.574.

Conclusion: Given the correlation results obtained, the clinical experience of physicians will still have one of the leading roles in diagnosing acute appendicitis in children.

Key Words: Appendicitis, Children, Leukocytes, Segmented neutrophils.

*Please cite this article as: Baskovic M, Zupancic B, Dominkovic A, Cizmic A, Jezek D. Correlation of Leukocyte Count and Percentage of Segmented Neutrophils with Pathohistological Findings of Appendix in Children. Int J Pediatr 2018; 6(1): 6851-59. DOI: [10.22038/ijp.2017.26748.2304](https://doi.org/10.22038/ijp.2017.26748.2304)

*Corresponding Author:

Marko Baskovic, MD, Department of Pediatric Surgery, Children's Hospital Zagreb, Klaićeva 16, Zagreb 10000, Croatia.

Email: baskovic.marko@gmail.com

Received date: Nov.17, 2017; Accepted date: Dec.22, 2017

1- INTRODUCTION

Appendicitis is the most common indication for an emergency operation in children's age. In everyday clinical practice diagnosis of appendicitis is usually based on clinical picture, physical examination, laboratory findings and ultrasound examination. Although none of the laboratory values has not high sensitivity and specificity for the diagnosis of appendicitis, leukocyte count and the percentage of segmented neutrophils are most commonly used. The increase in the percentage of segmented neutrophils is strongly associated with appendicitis, as only 3.7% of children without left shift have appendicitis (1).

These two lab markers are the most discriminating factor for appendicitis within 24 hours of the onset of pain. In order to objectively evaluate the clinical findings, in Alvarado score leukocytosis ($\geq 10 \times 10^9 / L$) with segmented neutrophils ($\geq 75\%$) carry 3 points while in Pediatric Appendicitis score carry 2 out of 10 points (2, 3). The aim of this study was to determine whether there is a statistically significant correlation between leukocyte count and the percentage of segmented neutrophils compared to the pathohistological finding of appendix in children.

2- MATERIALS AND METHODS

In order to ascertain whether there was a statistically significant correlation between leukocyte count and the proportion of segmented neutrophils compared to the pathohistological finding of children whose appendix has been removed, we retrospectively analyzed the data from the Hospital Information System and hospital documentation in the period from 1 January 2016 to 31 December 2016 (from the Children's Hospital Zagreb, Croatia). During this period 243 children were operated with a classic or laparoscopic approach. From the study, 25 patients were

excluded due to incomplete documentation, and 7 patients with periappendiceal abscess who subsequently went to elective appendectomy. The analysis was made on 211 patients. All analyzed laboratory findings (leukocyte count, percentage of segmented neutrophils) were made within 12 hours before surgery. The results of the pathohistological analysis are classified into five categories: normal findings, acute appendicitis, acute phlegmonous appendicitis, acute gangrenous appendicitis and acute gangrenous perforated appendicitis. The data obtained is displayed via the scatter diagram in Microsoft Excel. In order to ascertain whether there is a correlation between leukocyte count and the segmented neutrophils in relation to the pathohistological finding, Spearman correlation coefficient is calculated ($r_s = 1 - 6 \times \sum_{i=1...N} D_i^2 / [N(N^2-1)]$), the result of which is interpreted according to the following value ranges (0 - $\pm 0.25 \leftrightarrow$ no correlation, $\pm 0.26 - \pm 0.50 \leftrightarrow$ weak correlation, $\pm 0.51 - \pm 0.75 \leftrightarrow$ good correlation, $\pm 0.76 - \pm 1 \leftrightarrow$ excellent correlation). Results were interpreted if they were statistically significant ($p < 0.05$). We determined, based on our results, the specificity and sensitivity of leukocyte count and the percentage of segmented neutrophils used in the calculation of Alvarado and Pediatric Appendicitis score. We calculated area under the curve for leukocytes and segmented neutrophils. The analysis was made in XLSTAT (statistical *software* for Microsoft Excel).

3- RESULTS

Out of a total of 211 patients (boys - 118 (55.9%); girls - 93 (44.1%)), the youngest operated patient was 2 years and 3 months, while the oldest was 17 years and 9 months. The average age was 11 years and 5 months. The results of the pathohistological analysis were as follows:

normal finding - 30, acute appendicitis - 47, acute phlegmonous appendicitis - 79, acute gangrenous appendicitis - 43, acute gangrenous perforated appendicitis - 12. In the scatter diagrams (**Figure.1**, **Figure.2**), results of leukocyte count and the percentage of segmented neutrophils relative to the pathohistological finding are shown. One hundred and ninety (90%) patients had a leukocyte count greater than $10 \times 10^9/L$, while 144 patients (68.2%) had a percentage of segmented neutrophils greater than 75%. The results are presented in the contingency tables (**Table.1**, **Table.2**), with sensitivity for leukocyte count of 93% and specificity of 30%, while sensitivity for percentage of segmented neutrophils is 71% and specificity of 50% (**Figure.3**, **Figure.4**).

The Receiver operating characteristic (ROC) curve analysis for leukocyte count shows area under the curve of 0.648, while for segmented neutrophils of 0.574 (**Figure.5**, **Figure.6**).

By adding the leukocyte and pathohistological findings data into the Spearman coefficient correlation formula, $r_s = 0.29$ ($p = 3.61 \times 10^{-8}$) is obtained. For leukocytes, the correlation is weak. By adding the data for the segmented neutrophils into the formula for the Spearman coefficient of correlation, $r_s = 0.18$ ($p = 7.08 \times 10^{-5}$) is obtained. Correlation between the percentage of segmented neutrophils and pathohistological findings does not exist.




Fig.1: Scatter plot: relationship between pathohistological finding and leukocyte count.




Fig.2: Scatter plot: relationship between pathohistological finding and the percentage of segmented neutrophils.

Table-1: Contingency table: The relationship between pathohistological finding and leukocyte count.

Leukocyte count	Positive pathohistological findings	Negative pathohistological findings	Total
Leukocytes $\geq 10 \times 10^9/L$	169	21	190
Leukocytes $\leq 10 \times 10^9/L$	12	9	21
Total	181	30	211

Table-2: Contingency table: The relationship between pathohistological finding and the percentage of segmented neutrophils.

Percentage of segmented neutrophils	Positive pathohistological findings	Negative pathohistological findings	Total
Segmented neutrophils $\geq 75\%$	129	15	144
Segmented neutrophils $\leq 75\%$	52	15	67
Total	181	30	211




Fig.3: Sensitivity / Specificity for leukocytes.




Fig.4: Sensitivity / Specificity for segmented neutrophils.




Fig.5: ROC curve for leukocytes.




Fig.6: ROC curve for segmented neutrophils.

4- DISCUSSION

In our study we had a negative rate of appendectomy of 14.2% which is in accordance with the rates of negative appendectomy (NA) of 5% to 15% (4). The most recent reports quote NA rates below 5% (5, 6), which we should strive in the future. Bates et al. analyzed all NAs, to detect common characteristics that, in the future, might further decrease false-positive rate (7). Absence of an elevated white blood cell count (WBC) count is a risk factor for NA. Withholding appendectomy for WBC counts, 9,000 and 8,000 per mL reduces the NA rate to 0.6% and 1.2%. In our study, out of a total of 30 negative appendectomy, 9 patients (30%) had an orderly leukocyte count ($\leq 10 \times 10^9/L$), while 15 patients (50%) had a percentage of segmented neutrophils $< 75\%$. Stefanutti et al. concluded that WBC count at admission has no proven additional value in the diagnosis of acute appendicitis and can be omitted without modifying diagnostic pathway and without affecting diagnostic accuracy (8).

The purpose of Mathews et al. study was to determine if Immature Granulocytes (IG)% would add additional discriminatory ability, in conjunction with traditionally utilized laboratory values, in differentiation of acute and perforated appendicitis in a pediatric population. In that study, they found that elevated C-reactive protein (CRP) and presence of a left shift were reliable predictors of perforated appendicitis. However, IG% did not add any additional benefit in making the diagnosis (9). In the Kulik et al. study, the pediatric Appendicitis Score (PAS) and Alvarado scores were the best validated but neither met the current performance benchmarks. A high quality well validated and consistently high-performing clinical prediction rules (CPR) was not identified. They concluded that further research is needed before a CPR for children with suspected appendicitis can be used in

routine practice (10). According to Wang et al. study (11), elevated WBC counts had a sensitivity of 67% and a specificity of 80%. Using left shift alone as an indicator for appendicitis was associated with a sensitivity of 59% and a specificity of 90%. However, when a high WBC count and left shift were combined, the sensitivity climbed to 80%, and specificity remained at 79%. The sensitivity fell to 47% when both a high WBC count and left shift were analyzed, and specificity climbed to 94%. The positive likelihood ratio for a high WBC count and left shift was 9.8. Sensitivity of leukocytosis (93%) and shift to left (71%) in Gwynn study (12) was analogous to ours, while the specificity of leukocytosis was 38% and shift to left 68%. In the study Ko et al. (13) leukocytosis and shift-to-the-left were better predictors than CRP (the sensitivity and specificity are: leukocytosis ($> 10,000/mm^3$): 85.2%, 65%; leukocytosis with a shift-to-the-left (neutrophil $> 75\%$): 81.5%, 70%; elevated CRP (> 0.9 mg/dl): 70.4%, 65%). In our study, we did not calculate sensitivity and specificity for CRP because it is only available in only 34 patients.

We did not find an article that would show whether there was a statistically significant correlation between the leukocyte count and the percentage of segmented neutrophils compared to the pathohistological finding in children. Tatar et al. (14) calculated the correlation between appendiceal diameter - WBC count and appendiceal diameter - Alvarado score. The correlation between appendiceal diameter and WBC count was 80% ($p = 0.01 < 0.05$). The correlation between appendiceal diameter and Alvarado score was 78.7% ($p = 0.01 < 0.05$). Sousa-Rodrigues et al. (15) evaluated the possible correlation between the scale of Alvarado (EA) and macroscopic appearance (MA) of the appendix in patients with acute

appendicitis. The Spearman correlation test used for EA and MA was + 0.77 (95% CI 0.65-0.85, $p < 0.0001$). Bhatti Hafeez et al. (16) wanted to determine the correlation of WBC counts, age and duration of symptoms with severity of acute appendicitis. They determined that WBC counts and duration of symptoms are not good predictors of severity of disease in appendicitis. In conclusion of the study Grönroos et al. (17) WBC remained the best laboratory method for diagnosing uncomplicated acute appendicitis and seemed to be a very early marker of appendiceal inflammation. They calculated Spearman rank-order correlation between WBC, CRP, and lipoprotein-associated phospholipase A2 (Lp-PLA2).

Grönroos in his study (18) concluded that, contrary to adult patients, normal leucocyte count and CRP value do not effectively exclude acute appendicitis in children. Gulzar et al. (19) said that clinical assessment is the best criteria to reach a confident diagnosis. Total leucocyte count and other investigations should be used as diagnostic aid in doubtful cases but they don't replace the clinical skills of surgeons. In their study total leucocyte count was less than 10,000/mm³ in 38.8% cases and more than 10,000/mm³ in 61.2% cases. The sensitivity and specificity of raised TLC for acute appendicitis were 80% and 67%, respectively.

5- CONCLUSION

Although daily attempts to find better (more sensitive and more specific) tests in the diagnosis of acute appendicitis in children, leukocyte count and the percentage of segmented neutrophils within the Alvarado score continue to be the leading position with the highest sensitivity and specificity of all tests. Given the correlation results obtained, the clinical experience of physicians will still

have one of the leading roles in diagnosing acute appendicitis in children.

6- CONFLICT OF INTEREST: None.

7- REFERENCES

1. Rentea RM, St Peter SD. Pediatric Appendicitis. *Surg Clin North Am.* 2017; 97(1):93-112.
2. Alvarado A. A practical score for the early diagnosis of acute appendicitis. *Ann Emerg Med.* 1986; 15(5):557-64.
3. Samuel M. Pediatric appendicitis score. *J Pediatr Surg.* 2002; 37(6):877-81.
4. Buchman TG, Zuidema GD. Reasons for delay of the diagnosis of acute appendicitis. *Surg Gynecol Obstet.* 1984; 158(3):260-66.
5. Bachur RG, Hennelly K, Callahan MJ, Chen C, Monuteaux MC. Diagnostic imaging and negative appendectomy rates in children: effects of age and gender. *Pediatrics.* 2012; 129(5):877-84.
6. Oyetunji TA, Ong'uti SK, Bolorunduro OB, Cornwell EE III, Nwomeh BC. Pediatric negative appendectomy rate: trend, predictors and differentials. *J Surg Res.* 2012; 173(1):16-20.
7. Bates MF, Khander A, Steigman SA, Tracy TF Jr, Luks FI. Use of white blood cell count and negative appendectomy rate. *Pediatrics.* 2014; 133(1):39-44.
8. Stefanutti G, Sabatti M, Gobbi D, Ghirardo V, Gamba PG. Values of white blood cell count in the diagnosis of acute appendicitis. *Pediatr Med Chir.* 2002; 24(5):368-73.
9. Mathews EK, Griffin RL, Mortellaro V, Beierle EA, Harmon CM, Chen MK, Russell RT. Utility of immature granulocyte percentage in pediatric appendicitis. *J Surg Res.* 2014; 190(1):230-4.
10. Kulik DM, Uleryk EM, Maguire JL. Does this child have appendicitis? A systematic review of clinical prediction rules for children with acute abdominal pain. *J Clin Epidemiol.* 2013; 66(1):95-104.

11. Wang LT, Prentiss KA, Simon JZ, Doody DP, Ryan DP. The use of white blood cell count and left shift in the diagnosis of appendicitis in children. *Pediatr Emerg Care.* 2007; 23(2):69-76.
12. Gwynn LK. The diagnosis of acute appendicitis: clinical assessment versus computed tomography evaluation. *J Emerg Med.* 2001; 21(2):119-23.
13. Ko YS, Lin LH, Chen DF. Laboratory aid and ultrasonography in the diagnosis of appendicitis in children. *Zhonghua Min Guo Xiao Er Ke Yi Xue Hui Za Zhi.* 1995; 36(6):415-9.
14. Gunes Tatar I, Yilmaz KB, Sahin A, Aydin H, Akinci M, Hekimoglu B. Evaluation of Clinical Alvarado Scoring System and CT Criteria in the Diagnosis of Acute Appendicitis. *Radiology Research and Practice.* 2016; 9739385.
15. Sousa-Rodrigues CF, Rocha, AC, Rodrigues AKB, Barbosa FT, Ramos FWS, Valões SHC. Correlation between the Alvarado Scale and the macroscopic aspect of the appendix in patients with appendicitis. *Revista do Colégio Brasileiro de Cirurgiões.* 2014; 41(5): 336-339.
16. Bhatti Hafeez AB, Dawood A, Memon F, Zaman J. Acute Appendicitis: Can WBC count, age and duration of symptoms predict severity of disease. *Pakistan Journal of Surgery.* 2009; 25(3): 167-70.
17. Grönroos JM, Forsström JJ, Irjala K, Nevalainen TJ. Phospholipase A2, C-reactive protein, and white blood cell count in the diagnosis of acute appendicitis. *Clin Chem.* 1994;40(9):1757-60.
18. Grönroos JM. Do normal leucocyte count and C-reactive protein value exclude acute appendicitis in children? *Acta Paediatr.* 2001; 90(6): 649-51
19. Gulzar S, Umar S, Dar GM, Rasheed R. Acute appendicitis - importance of clinical examination in making a confident diagnosis. *Pak J Med Sci.* 2005; 21: 125-32.